

## AIDS Diagnoses by Top 10 States/Dependent Areas

The 10 states or dependent areas reporting the highest number of AIDS diagnoses in 2008 were:

State/Dependent Area	# of AIDS Diagnoses, 2008
California	4,835
Florida	4,766
New York	4,571
Texas	2,924
Georgia	1,908
Maryland	1,557
New Jersey	1,527
Pennsylvania	1,402
Illinois	1,305
North Carolina	1,157

State/Dependent Area	# of Cumulative AIDS Diagnoses Through 2008*		
	Adults or Adolescents	Children (<13)	Total
New York	190,363	2,390	192,753
California	159,606	687	160,293
Florida	116,041	1,571	117,612
Texas	76,674	396	77,070
New Jersey	53,756	801	54,557
Georgia	38,054	245	38,300
Pennsylvania	37,842	375	38,217
Illinois	37,592	288	37,880
Maryland	35,395	330	35,725
Puerto Rico	32,057	406	32,463

\* From the beginning of the epidemic through 2008.

[Go to top](#)

## Persons Living with an AIDS Diagnosis

At the end of 2007, the **estimated number** of persons living with an AIDS diagnosis in the United States and dependent areas was **470,902**. In the 50 states and the District of Columbia, this included **458,686** adults and adolescents, and **908** children under age 13 years.

Estimated numbers resulted from statistical adjustment that accounted for delays in reporting to the health department (but not for incomplete reporting) and missing risk factor information, where appropriate.

Because of delays in reporting of deaths, data are only available through the end of 2007. The exclusion of data from the most recent year allows at least 18 months for deaths to be reported and for these persons to be removed from calculations of persons living with an AIDS diagnosis.

Totals include persons of unknown race/ethnicity. Because totals for the estimated numbers were calculated independently of the values for the subpopulations, the subpopulation values may not equal the totals.

[Go to top](#)

### **Deaths of Persons with an AIDS Diagnosis**

In 2007, the **estimated number** of deaths of persons with an AIDS diagnosis in the United States and dependent areas was **18,089**. In the 50 states and the District of Columbia, this included **17,613** adults and adolescents, and **6** children under age 13 years.

The cumulative **estimated number** of deaths of persons with an AIDS diagnosis in the United States and dependent areas, through 2007, was **597,499**. In the 50 states and the District of Columbia, this included **571,453** adults and adolescents, and **4,931** children under age 13 years.

Deaths of persons with an AIDS diagnosis may be due to any cause.

Estimated numbers resulted from statistical adjustment that accounted for delays in reporting to the health department (but not for incomplete reporting) and missing risk factor information, where appropriate. Because of delays in reporting of deaths, data are only available through the end of 2007. The exclusion of data from the most recent year allows at least 18 months for deaths of persons with an AIDS diagnosis to be reported.

Totals include persons of unknown race/ethnicity. Because totals for the estimated numbers were calculated independently of the values for the subpopulations, the subpopulation values may not equal the totals.

[Go to top](#)

### **State-by-State HIV infection and AIDS Data**

- [NCHHSTP State Profiles](#)
- [Statehealthfacts.org](#) ■ provides state-by-state information about new and cumulative AIDS diagnoses, AIDS diagnosis rates, persons living with an AIDS

diagnosis, AIDS deaths, diagnoses of HIV infections, HIV testing statistics and policies, additional AIDS-related state policies, Ryan White funding and funding for HIV prevention, and AIDS Drug Assistance Programs, including budget, client, and expenditure data from the Kaiser Family Foundation.

[Go to top](#)

[Go to top](#)

Last Modified: July 27, 2010

Last Reviewed: July 27, 2010

Content Source:

[Divisions of HIV/AIDS Prevention](#)

[National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention](#)



<http://www.cdc.gov/hiv/topics/surveillance/basic.htm#area>

\